



35 Waverly Place, New York 5

# AEC Computing and Applied Mathematics Center

## AEC RESEARCH AND DEVELOPMENT REPORT

NYO-2544

SHIMWOOD PROGRESS REPORT

No. 3

(July 1958 to July 1959)

H. Grad and J. Berkowitz


July 31, 1959

## Institute of Mathematical Sciences

NEW YORK UNIVERSITY

NEW YORK, NEW YORK

140-2544



This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, express or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.



As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

UNCLASSIFIED

AEC COMPUTING AND APPLIED MATHEMATICS CENTER  
Institute of Mathematical Sciences  
New York University

NYO-2544

SHERWOOD PROGRESS REPORT

No. 3

(July 1958 to July 1959)

H. Grad and J. Berkowitz

July 31, 1959

Contract No. AT(30-1)-1480

- 1 -

UNCLASSIFIED

## Personnel

The average number of equivalent full-time scientific staff members was about 16 theoreticians and 2 experimenters. The total number of individuals involved was about 25; these include 13 of faculty rank or with Ph.D.'s (8.5 equivalent full-time) and 12 research assistants (9.5 equivalent full-time).

## Propagation of Hydromagnetic Waves

The general problem is the description of the properties of small amplitude signals propagating in a plasma. To this end, the computation of a dispersion relation should be considered to be the beginning rather than the end of the problem. In specific cases which could be evaluated in detail, the properties of plane waves were found to be not only incomplete but even misleading. For example, non-plane fronts can be found which propagate faster in some directions than the maximum speed of propagation of a plane wave in any direction [22-VIII]; similarly, one mode whose plane wave fronts propagate slower than those of another mode in every direction can catch up and pass the second mode if the front is not plane [3]. Complete modes (as distinguished from plane waves) are found to separate only in special cases and, even then, only in a certain subtle sense [3].

The types of question which were approached are the possibility of separation of modes (in arbitrary disturbances rather than plane waves), channeling of energy propagation and radiation patterns, and the general propagation of arbitrary wave fronts including refraction, focussing, etc. with the idea of eventual application to localized plasma diagnostics. The deeper questions force one to study, at least initially, some of the simpler models of plasma behavior. For the classical perfectly conducting fluid, an extensive and almost complete analysis has been made of the propagation of arbitrary

wave fronts into a general inhomogeneous medium (geometrical optics)[18]. Not only the position of the front, but also the variation in amplitude are given by simple formulas. The more difficult problem of general energy and signal transfer (not necessarily related to a wave front) has been successfully attacked in special geometries [3] and is being extended to more general cases by the use of recently refined asymptotic techniques [19]. For perturbations about a constant magnetic field, one mode was found which propagates purely one-dimensionally, guided along the magnetic lines [3]. Conceivably, this could be used as a diagnostic tool to obtain average properties of an open-ended plasma along a single magnetic line. The problem of propagation into a non-uniform field is being studied.

Extension of these results to more realistic plasma models using the Boltzmann equation are under way. Since many of the questions have never been satisfactorily answered for even a simple gas without ionization or electromagnetic effects present, it was found to be necessary to first develop some parts of this simpler theory before dealing with plasmas. As a first step, an asymptotic approximation was obtained to the time-independent fundamental solution of the Boltzmann equation [4]. This is being extended to time-dependent propagation and to similar analysis in plasmas.

## Particle Orbits

The proof of the asymptotic validity of the guiding center expansion [27] was submitted for publication [10].

A large number of general results concerning adiabatic invariants and the asymptotic behavior of charged particle orbits was obtained. In September (this was announced informally at the Geneva conference) C. S. Gardner proved the existence of the "second" adiabatic invariant (even to arbitrarily high order) for axially symmetric systems. These results were extended [40a], using classical perturbation theory of Hamiltonian systems to obtain

- 1) a new and simplified proof of the invariance of the magnetic moment to arbitrarily high order
- 2) under appropriate circumstances (roughly some sort of mirror arrangement -- but not necessarily axially symmetric) the invariance to arbitrary order of a second adiabatic invariant.
- 3) under appropriate circumstances the invariance of a third invariant
- 4) extension of all these results to relativistic particles.

The Hamiltonian treatment yields a reduction of the original sixth order system to fourth order on elimination of the magnetic moment as a "constant" of the motion and a similar reduction to second order on elimination of the second invariant. What remains is a system of one degree of freedom

describing the motion of the "line" to which the particle is "tied". A brief account was submitted for publication [13], and a more detailed version is being prepared for publication.

These adiabatic results do not give much information about deviations from adiabaticity. Some preliminary analytic results in this direction were obtained, but much more work is required.

Special problems in which applications of particle orbit theory arise are described under Shock Theory, Equilibrium and Stability, and Cusped Geometries.

### Shock Theory

The analysis of the self-consistent steady shock (non-colliding particles) propagating in a direction perpendicular to the magnetic field was continued. It had previously been observed that there are at least two significantly different sources of entropy production, in particular one originating from particles which are reflected back by their first encounter with the charge separation electric field [1c]. It was now observed that there are parameter values for which this entropy increase is an appreciable part of the maximum which could occur across a complete shock [40c]. Thus, even ignoring other mechanisms, there can be appreciable irreversible heating of the ions in a very thin layer. Specifically,



there are interesting ranges of the physical parameters which yield an essentially discontinuous shock front across which ions alone are irreversibly heated; this front is followed by an approximately adiabatic periodic wave train. A full-scale analysis and computation of the magnitude of this heating as a function of various parameters was carried out and will be issued shortly [11].

Details of some of the earlier work in this subject were issued in [2].

Several fluid-type approximations to this problem were continued (cf. [35g]), but, thus far, unsuccessfully (i.e., in prediction of a damped shock profile).

### Transport Theory

By making strong use of symmetry relations, it was found to be possible to elucidate and clarify the jungle of transport coefficients and collision integrals which arise from the Boltzmann equation [14]. The program aimed at improved values for these coefficients, see [29] and [7], was continued. A dictionary of collision integrals is being prepared (L. Sirovich) to avoid future repetitions of the extensive redundant calculations in the literature.

The question of the slowing down and thermalization of a single fast ion was treated, [40d], [12]. Care had to be taken to include both electron and ion effects since either one dominated in different regions. Quite complicated thermalization patterns were found, including development of a secondary peak in the distribution and the transition from an initial  $\delta$ -function through a distribution much wider than the ultimate Maxwellian before final thermalization. The final decay to Maxwellian was found to be approximately exponential, but the decay time depends strongly on the initial energy of the ion. In other words, reference to a universal ion "thermalization" time can be quite misleading.

The asymptotic evaluation of the tail end of the ion distribution and resultant correction in the thermonuclear reaction rate [28], [1e], was improved to obtain an exact rather than an asymptotic result (R. Liboff - numerical results not yet complete).

Two questions of anomalous resistivity induced by plasma oscillations were considered. The first depends on the increased dispersion of a particle due to the presence of externally induced plasma oscillations which are only slowly damped [40b]. This effect can be exceptionally large in a sheath. The second results from the introduction of plasma oscillations into the formulation of Ohm's law, thereby modifying the diffusion rate of plasma into field (J. Thurber, not yet complete). These

two effects are independent; in particular, the plasma resistivity value to be used in the second problem should be the corrected value as given by the first.

### Equilibrium and Stability

The problem of the self-consistent equilibrium of a mirror-machine geometry in the guiding center limit (with non-diagonal stress tensor, generalizing the fluid model,  $\nabla p = J \times B$ ) was solved in certain limiting cases (A. Oppenheim - report to appear). More general solutions also seem to be accessible and are being studied. A stability analysis will follow.

Existence theorems have been proved for a variety of axially symmetric and toroidal free boundary equilibria [40e] (M. Schechter and C. Morawetz - detailed report to appear).

The problem of the stability of a toroidal "stabilized" pinch has been formulated and prepared for computation (R. D. Richtmyer, R. Lüst, B. Suydam).

A stability analysis using the guiding center fluid equations (with non-diagonal stress tensor) [34h] was found to yield instability when the ratio of parallel to perpendicular pressure is either too large or too small [40f]. At

second glance, this result is found to be illusory or, at least, hard to interpret since the equations of motion are found to be not well-posed when the stated inequalities are exceeded. In other words, the motion is not unstable; it is nonexistent. This result may cast doubt on all stability analyses which use the guiding center limit since it is this limit which seems to produce the paradoxical result.

Comprehensive and detailed reports covering the material which was briefly summarized in the two Geneva papers [1b] and [1d] will be issued shortly, [8-IX] and [8-X].

### Cusped Geometries

Work in this subject consisted mainly in the continuation of problems which have been under study for some time. These include the computation of axially symmetric free boundary configurations and the computation of nonadiabatic orbits in cusped magnetic fields for the improved estimation of particle loss rates.

References: [1a], [25], [6], [9].

### Experimental

The experimental program was terminated as of July 1959. A final report is being prepared [15].

## BIBLIOGRAPHY

ISSUED JULY 1958 - JULY 1959

- [1] Papers presented at Geneva, September 1958, NYO-2538  
This report contains corrected manuscripts of the following five papers:
  - a. Cusped Geometries by J. Berkowitz, K. O. Friedrichs, H. Goertzel, H. Grad, J. Killeen, E. Rubin
  - b. Hydromagnetic Equilibria and Force-Free Fields by H. Grad and H. Rubin
  - c. Hydromagnetic Shock Waves in High-Temperature Plasmas C. S. Gardner, H. Goertzel, H. Grad, C. S. Morawetz, M. H. Rose, H. Rubin
  - d. Problems in Magnetohydrodynamic Stability by J. Berkowitz, H. Grad and H. Rubin
  - e. Thermonuclear Reaction Rate in a Discharge by H. Grad
- [2] Magneto-hydrodynamic Shock Structure Using Friction. Cathleen S. Morawetz, NYO-8677, January 12, 1959.
- [3] Propagation of Magnetohydrodynamic Waves without Radial Attenuation. Harold Grad, NYO-2537, January 15, 1959.  
Presented at the Third Lockheed Symposium on Magneto-hydrodynamics on November 21, 1958, also at APS Fluid Dynamics Division Colloquium on May 2, 1959.  
Also to be issued in the Proceedings of the Symposium.
- [4] Equations of Flow in a Rarefied Atmosphere. Harold Grad, NYO-2543, June 17, 1959.  
Presented at the Symposium on the Aerodynamics of the Upper Atmosphere at the Rand Corp. on June 8-10, 1959.  
Also issued in the Proceedings of the Symposium.
- [5] Principles of the Kinetic Theory of Gases. Harold Grad, Handbuch der Physik, Vol. XII, p. 205.
- [6] Theory of Cusped Geometries II. Particle Losses. Jerome Berkowitz, NYO-2536, January 6, 1959.

- [7] Transport Coefficients Determined Using the Shielded Coulomb Potential. Richard L. Liboff, Physics of Fluids, Vol. 2, January-February, 1959, p. 40.

TO APPEAR SHORTLY

- [8] Notes on Magnetohydrodynamics. NYO-6486  
IX Motion and Stability of Hydromagnetic Free Surfaces.  
A. A. Blank and H. Grad  
X Hydromagnetic Equilibria and Force-Free Fields.  
H. Grad and H. Rubin
- [9] Cusped Geometries III. Particle Losses with Crossed Fields. John Killeen
- [10] On the Asymptotic Series Expansion of the Motion of Charged Particle in Slowly-Varying Fields. J. Berkowitz and C. S. Gardner, Comm. Pure & Appl. Math., Vol. XII, No. 3, August 1959.
- [11] Magnetohydrodynamic Shock Structure Without Collisions. Cathleen S. Morawetz
- [12] Thermalization of a Fast Ion in a Plasma. Herbert C. Kranzer
- [13] Remarks on Adiabatic Invariance. C. S. Gardner, submitted to Phys. Rev.
- [14] Theoretical Work in the Kinetic Theory of Gases. Harold Grad, Symposium on Superaerodynamics and Rarefied Gas Aerodynamics, Nice, July, 1958.

- [15] Experiments on Supersonic Plasma Flow Along Magnetic Fields. II. D. Wetstone, NYO-2877, July 31, 1959.

OTHER IMS REPORTS ON SUBJECTS RELATED TO SHERWOOD

- [16] Resolution of an Initial Shear-Flow Discontinuity in One-Dimensional Hydromagnetic Flow. J. Bazer, Astrophysical Journal, Vol. 128, No. 3, November, 1958, p. 686.
- [17] Hydromagnetic Shocks. J. Bazer and W. B. Ericson. Astrophysical Journal, Vol. 129, No. 3, May 1959, p. 758.
- [18] Propagation of Weak Hydromagnetic Discontinuities. Jack Bazer and Owen Fleischman, Research Report No. MH-10, Div. of EM Research, Inst. of Math. Sci., NYU, March 1959, also Physics of Fluids, Vol. 2, No. 4, July-August 1959, p. 366.
- [19] Exact and Asymptotic Solutions of the Cauchy Problem. D. Ludwig
- [20] Symmetric Positive Linear Differential Equations. K. O. Friedrichs, Comm. Pure & Appl. Math., Vol. XI, No. 3, August 1958, p. 333.

PREVIOUSLY ISSUED

- [21] Notes on Magnetohydrodynamics. First series, issued 1954. No official distribution. Replaced by references [22 I, III, IV, VII].
- [22] Notes on Magnetohydrodynamics. NYO-6486
- I. General Fluid Equations. Harold Grad, August 1956.
  - II. Dimensional Considerations. Harold Grad and Marian H. Rose, August 1956.
  - III. Special Solutions. Harold Grad, August 1956.
  - IV. Ohm's Law. Harold Grad, August 1956.
  - V. Theory of Maxwell's Equations without Displacement Current. A. A. Blank, K. O. Friedrichs, H. Grad, November 1957.
  - VI. Fluid Magnetic Equations -- General Properties. A. A. Blank and Harold Grad, July 1, 1958.
  - VII. Fluid Dynamical Analogies. A. A. Blank and Harold Grad, July 15, 1958.
  - VIII. Nonlinear Wave Motion. K. O. Friedrichs and H. Kranzer, July 31, 1958. Originally issued as: Nonlinear Wave Motion in Magnetohydrodynamics. K. O. Friedrichs, LAMS-2105, Los Alamos Scientific Laboratory. Written September 1954, issued 1957.
- [23] On the Diffusion of a Conducting Fluid Across a Magnetic Field. Marian H. Rose, NYO-7691, July 1, 1956.
- [24] On the Structure of a Steady Hydromagnetic Shock -- One-Fluid Theory. Marian H. Rose, NYO-7693, October 1956.
- [25] The Theory of Cusped Geometries I. General Survey. Harold Grad, NYO-7969, December 1, 1957 (Reissued January 20, 1959).
- [26] Basic Notions of Relativistic Hydromagnetics. Paul Reichel, NYO-7697, April 15, 1958.



- [27] On the Asymptotic Series Expansion of the Motion of a Charged Particle in Slowly-Varying Fields. J. Berkowitz and C. S. Gardner, NYO-7975, December 23, 1957.
- [28] Thermonuclear Reaction Rates in an Electrical Discharge. Harold Grad, NYO-7977, January 23, 1958.
- [29] Moment Description of Gas Mixtures - I. Ignace I. Kolodner, NYO-7980, September 10, 1957.
- [30] Experiments on Supersonic Plasma Flow Along Magnetic Fields. D. Finkelstein, M. Earlich, I. Livingston, and D. Wetstone, NYO-7981, March 27, 1958.
- [31] Computations of the Transport Coefficients Using the Shielded Coulomb Potential. Richard L. Liboff, NYO-8669, June 20, 1958.

PAPERS PRESENTED AT SHERWOOD MEETINGS

- [32] Princeton, October 1954. WASH-184, issued January 1955.
  - a. Talk by Harold Grad, p. 144.
  - b. Talk by K. O. Friedrichs, p. 148.
- [33] UCRL, February 1955. WASH-289, issued June 1955.
  - a. Survey of Activities. Harold Grad, p. 108.
  - b. On the Diffusion of Conducting Fluids in Magnetic Fields. Marian H. Rose, p. 111.
  - c. Motion and Stability of Free Surfaces. Harold Grad, p. 115.
  - d. Ohm's Law and Superconductivity. Harold Grad, p. 124.
- [34] Princeton, October 1955. TID-7503, issued February 1956.
  - a. Stability of Free Surfaces. Harold Grad, p. 238.

- b. Dynamical Stabilization of the Pinch: Preliminary Report. Jerome Berkowitz, p. 241.
- c. Local Stability of an Incompressible Plasma Mixed with a Magnetic Field. Hanan Rubin, p. 247.
- d. Theory of the Cuspoid. Harold Grad, p. 319.
- e. Projection of Plasma Across a Magnetic Field. W. H. Bostick, D. R. Lasher, David Finkelstein, p. 355.
- f. On the Structure of a shock in a Magnetic Field. Marian H. Rose, p. 475.
- g. Motion of a Particle in a Uniform Magnetic Field. John S. Lomont, p. 480.
- h. A Guiding Center Fluid. Harold Grad, p. 495.

[35] Gatlinburg, June 1956. TID-7520, issued September 1956.

- a. Survey of Progress to Date in the NYU Program. Harold Grad, p. 95.
- b. Developments in the Theory of Cusped Geometries. Harold Grad, p. 148.
- c. Some New Developments in Stability of Free Surfaces. Harold Grad, p. 373.
- d. Computation of Two-Dimensional Free Cusped Surfaces. Jerome Berkowitz, p. 394.
- e. Computation of Three Dimensional Free Cusped Surfaces. Herbert Goertzel, p. 400.
- f. General Adiabatic Theory. Harold Grad, p. 402.
- g. On the Structure of a Steady Hydromagnetic Shock Wave. Marian H. Rose, p. 547.
- h. Velocities of Vacuum Spark Plasmas. David Finkelstein, p. 589.

[36] Berkeley, February 1957, TID-7536, issued September 1957.

- a. Survey of the NYU Program, H. Grad, Pt. 2, p. 40.
- b. Stability of Magnetic Mirror Machines, H. Rubin, Pt. 2, p. 193.

- c. Morphology of Stable Geometries, H. Grad, Pt. 2, p. 196.
  - d. Particle Losses in Crossed Fields, J. Killeen, Pt. 2, p. 200.
  - e. Rigorous Justification of Kruskal's Asymptotic Series, C. Gardner and J. Berkowitz, Pt. 1, p. 10.
  - f. Shock Heating, H. Grad, Pt. 1, p. 32.
  - g. Shock Profiles, C. Morawetz, Pt. 1, p. 36.
  - h. Vacuum Spark Plasmas, D. Finkelstein, G. Sawyer, T. Stratton, Pt. 2, p. 542
- [37] Harwell, June 1957. AERE-GP/R-2371, issued 1957.
- a. Survey of NYU Activities, H. Grad, Vol. 1, p. 2.1.
  - b. Recent Developments at NYU, H. Grad, Vol. 2, p. 15.1.
- [38] Washington, February 1958, TID-7558, issued October 1958.
- a. Justification of Debye Cut-Off and Some Quantum Considerations, R. Liboff, p. 342.
  - b. Steady State Shock Model, C. S. Morawetz, p. 311.
  - c. Qualitative Considerations Concerning Stability, H. Grad, p. 368.
  - d. Neutron Emission in a Discharge, H. Grad, p. 516.
  - e. Wave in Plasma in Magnetic Field Caused by Application of Crossed Electric Field, C. S. Gardner, p. 307.
  - f. Force Free Electromagnet inside a Spherical Shell, G. K. Morikawa, p. 428.
- [39] New York, May 1958.
- a. Steady State Shock Model, C. S. Morawetz.
  - b. Steady State Waves in a Plasma: Asymptotic Behavior at Infinity in One-Dimensional Rectilinear Geometry, J. Calkin.
  - c. Stability of Relativistic Plasma Streams, D. Finkelstein.
  - d. Stable Mirror Machines, H. Kranzer and H. Rubin.
  - e. Explicit Equilibrium Configurations, H. Rubin.

- f. Reciprocal and Conjugate Vectors Applied to Equilibrium and Force-Free Fields, H. Rubin.
- g. Collisional Particle Losses in a Homopolar Device, H. Grad.
- h. Transport Coefficients of a Plasma by the Method of Moments, I. I. Kolodner.

[40] Gatlinburg, April 1959.

- a. Adiabatic Invariants of Charged-Particle Motion, C. S. Gardner.
- b. Increased Dispersion and Resistivity in a Nonsteady Plasma, H. Grad.
- c. Magneto-Hydrodynamic Shock Structure without Collisions, C. S. Morawetz and H. Goertzel.
- d. Thermalization of a Fast Ion in a Plasma, H. C. Krieger.
- e. Some Axially Symmetric Problems in Magneto-Hydrodynamics, M. Schechter.
- f. On the Stability of a Homogeneous Plasma with Non-isotropic Pressure, R. Lust.

red

NYO  
2544

Grad

Sherwood Progress Report  
TITLE

No. 3

BIRROWER S NAME

ROOM  
NUMBER

~~from~~ ~~Dr~~ Lissie

New York 3, N. Y.  
4 Washington Place

[illegible]

